

THE RIEMANNIAN SECTIONAL CURVATURE OPERATOR OF THE WEIL-PETERSSON METRIC AND ITS APPLICATION

YUNHUI WU

ABSTRACT. Fix a number $g > 1$, let S be a close surface of genus g , and $\text{Teich}(S)$ be the Teichmüller space of S endowed with the Weil-Petersson metric. In this paper we show that the Riemannian sectional curvature operator of $\text{Teich}(S)$ is non-positive definite. As an application we show that any twist harmonic map from rank-one hyperbolic spaces $H_{Q,m} = Sp(m,1)/Sp(m) \cdot Sp(1)$ or $H_{O,2} = F_4^{-20}/SO(9)$ into $\text{Teich}(S)$ is a constant.

1. INTRODUCTION

Let S be a closed surface of genus g where $g > 1$, and T_g be the Teichmüller space of S . T_g carries various metrics that have respective properties. For example, the Teichmüller metric is a complete Finsler metric. The McMullen metric, Ricci metric, and perturbed Ricci metric have bounded geometry [16, 17, 18]. The Weil-Petersson metric is Kähler [1] and incomplete [5, 26]. There are also some other metrics on T_g like the Bergman metric, Caratheodory metric, Kähler-Einstein metric, Kobayashi metric, and so on. In [16, 17], the authors showed that some metrics listed above are comparable. In this paper we focus on the Weil-Petersson case. Throughout this paper, we let $\text{Teich}(S)$ denote T_g endowed with the Weil-Petersson metric. The geometry of the Weil-Petersson metric has been well studied in the past decades. One can refer to Wolpert's recent nice book [29] for details.

The curvature aspect of $\text{Teich}(S)$ is very interesting, which plays an important role in the geometry of Weil-Petersson metric. This aspect has been studied over the past several decades. Ahlfors in [1] showed that the holomorphic sectional curvatures are negative. Tromba [24] and Wolpert [27] independently showed the sectional curvature of $\text{Teich}(S)$ is negative. Moreover, in [27] the author proved the Royden's conjecture, which says that the holomorphic curvatures are bounded above by a negative number that only depends on the topology of the surface, by establishing the curvature formula (see theorem 2.4). Wolf in [25] used harmonic tools to give another proof of this curvature formula. After that, people has been applying this formula to study the curvature of $\text{Teich}(S)$ in more detail. For example, in [21] Schumacher showed that $\text{Teich}(S)$ has strongly negative curvature in the sense of Siu (see [22]) that is stronger than negative sectional curvature. Huang in [10] showed there is no negative upper bound for the sectional curvature. In [15] Liu-Sun-Yau also used Wolpert's curvature formula to show that $\text{Teich}(S)$ has dual Nakano negative curvature, which says that the complex curvature operator on the dual tangent bundle is positive in some sense. For some other related problems one can refer to [3, 10, 11, 16, 17, 23, 28, 30].

Let $X \in \text{Teich}(S)$. We can view X as a hyperbolic metric on S . One of our purposes in this paper is to study the Riemannian sectional curvature operator of

$\text{Teich}(S)$ at X . The method in this paper is highly influenced by the methods in [15, 21, 27], which essentially applied the curvature formula, the Cauchy-Schwarz inequality and the positivity of the Green function for the operator $(\Delta - 2)^{-1}$, where Δ is the Beltrami-Laplace operator on X . What we need more in this paper is the symmetry of the Green function for $(\Delta - 2)^{-1}$.

Before giving any statements let us state some necessary background. Let X be a point in $\text{Teich}(S)$, and $T_X \text{Teich}(S)$ be the tangent space that is identified with the harmonic Beltrami differentials at X . Assume that $\{\mu_i\}_{i=1}^{3g-3}$ is a basis for $T_X \text{Teich}(S)$, and $\frac{\partial}{\partial t_i}$ is the vector fields corresponding to μ_i . Locally, t_i is a holomorphic coordinate around X ; let $t_i = x_i + \mathbf{i}y_i$. $(x_1, x_2, \dots, x_{3g-3}, y_1, y_2, \dots, y_{3g-3})$ gives a real smooth coordinate around X . Since $\text{Teich}(S)$ is a Riemannian manifold, it is natural to define the curvature tensor on it, which is denoted by $R(\cdot, \cdot, \cdot, \cdot)$. Let $T \text{Teich}(S)$ be the real tangent bundle of $\text{Teich}(S)$ and $\wedge^2 T \text{Teich}(S)$ be the wedge product of two copies of $T \text{Teich}(S)$. The curvature operator Q is defined on $\wedge^2 T \text{Teich}(S)$ by $Q(V_1 \wedge V_2, V_3 \wedge V_4) = R(V_1, V_2, V_3, V_4)$ and extended linearly, where V_i are real vectors. It is easy to see that Q is a bilinear symmetric form (one can see more details in [12]).

Now we can state our first result.

Theorem 1.1. *Let $S = S_g$ be a closed surface of genus $g > 1$ and $\text{Teich}(S)$ be the Teichmüller space of S endowed with the Weil-Petersson metric. And let \mathbf{J} be the almost complex structure on $\text{Teich}(S)$ and Q be the curvature operator of $\text{Teich}(S)$. Then, for any $X \in \text{Teich}(S)$, we have*

- (1) Q is non-positive definite, i.e., $Q(A, A) \geq 0$ for all $A \in \wedge^2 T_X \text{Teich}(S)$.
- (2) $Q(A, A) = 0$ if and only if there exists an element B in $\wedge^2 T_X \text{Teich}(S)$ such that $A = B - \mathbf{J} \circ B$.

$\mathbf{J} \circ B$ is defined in section 4.

A direct corollary is that the sectional curvature of $\text{Teich}(S)$ is negative [1, 24, 27]. Normally a metric of negative curvature may not have non-positive definite curvature operator (see [2]).

In the second part of this paper we will study harmonic maps from certain rank-one spaces into $\text{Teich}(S)$. For harmonic maps, there are a lot of very beautiful results when the target is either a complete Riemannian manifold with non-positive curvature operator or a complete non-positive curved metric space (see [6, 7, 31]). In particular, if the domain is either the Quaternionic hyperbolic space or the Cayley plane, different rigid results for harmonic maps were established in [9, 13, 19]. For harmonic maps into $\text{Teich}(S)$, one can refer to the nice survey [8]. In this paper we establish the following rigid result.

Theorem 1.2. *Let Γ be a lattice in a semisimple Lie group G which is either $Sp(m, 1)$ or F_4^{-20} , and $\text{Mod}(S)$ be the mapping class group of $\text{Teich}(S)$. Then, any twist harmonic map f from G/Γ into $\text{Teich}(S)$ with respect to each homomorphism $\rho : \Gamma \rightarrow \text{Mod}(S)$ must be a constant.*

The twist map f with respect to ρ means that $f(\gamma \circ Y) = \rho(\gamma) \circ f(Y)$ for all $\gamma \in \Gamma$.

Plan of the paper. In section 2 we provide some necessary background and some basic properties for the operator $D = -2(\Delta - 2)^{-1}$. In section 3 we establish the curvature operator formulas on different subspaces of $\wedge^2 T_X \text{Teich}(S)$ and show that the curvature operator is negative definite or non-positive definite on these different subspaces. In section 4 we establish the curvature operator formula for Q on $\wedge^2 T_X \text{Teich}(S)$ to prove theorem 1.1. In section 5 we finish the proof of theorem 1.2.

Acknowledgments. This paper is part of the author's thesis. The author is greatly indebted to his advisor, Jeffrey Brock, for his consistent encouragement and support. He would like to thank George Daskalopoulos for introducing this problem to the author and for many suggestions and discussions to help finish this article. He also would like to thank Zheng Huang, Georg Schumacher, Mike Wolf, and Scott Wolpert for useful conversations and suggestions.

2. NOTATIONS AND PRELIMINARIES

2.1. Surfaces. Let S be a closed surface of genus $g \geq 2$, and M_{-1} denote the space of Riemannian metrics with constant curvature -1 , and $X = (S, \sigma|dz|^2)$ be a particular element of M_{-1} . Diff_0 , which is the group of diffeomorphisms isotopic to the identity, acts by pull-back on M_{-1} . The Teichmüller space of S T_g is defined by the quotient space

$$M_{-1}/\text{Diff}_0.$$

The Teichmüller space has a natural complex structure, and its holomorphic cotangent space $T_X^* T_g$ is identified with the *quadratic differentials* $Q(X) = \varphi(z)dz^2$ on X . The *Weil-Petersson metric* is the Hermitian metric on T_g arising from the the *Petersson scalar product*

$$\langle \varphi, \psi \rangle = \int_S \frac{\varphi \cdot \bar{\psi}}{\sigma^2} dz d\bar{z}$$

via duality. We will concern ourselves primarily with its Riemannian part g_{WP} . Throughout this paper, we denote the Teichmüller space endowed with the Weil-Petersson metric by $\text{Teich}(S)$.

Setting $D = -2(\Delta - 2)^{-1}$, where Δ is the Beltrami-Laplace operator on X , we have $D^{-1} = -\frac{1}{2}(\Delta - 2)$. The following property has been proved in a lot of literature; for completeness, we still state the proof here.

Proposition 2.1. *Let D be the operator above. Then*

- (1) *D is self-adjoint.*
- (2) *D is positive.*

Proof of (1). Let f and g be two real-valued smooth functions on X , and $u = Df$, $v = Dg$. Then

$$\begin{aligned} \int_S Df \cdot g dA &= \int_S u \cdot \left(-\frac{1}{2}(\Delta - 2)v\right) dA = -\frac{1}{2} \int_S u \cdot (\Delta - 2)v dA \\ &= -\frac{1}{2} \int_S v \cdot (\Delta - 2)u dA = \int_S Dg \cdot f dA, \end{aligned}$$

where the equality in the second row follows from the fact that Δ is self-adjoint on closed surfaces. For the case that f and g are complex-valued, one can prove it

through the real and imaginary parts by using the same argument.

Proof of (2): Let f be a real-valued smooth functions on X , and $u = Df$. Then

$$\begin{aligned} \int_S Df \cdot f dA &= \int_S u \cdot \left(-\frac{1}{2}(\Delta - 2)u\right) dA = -\frac{1}{2} \int_S (u \cdot (\Delta u) - 2u^2) dA \\ &= \frac{1}{2} \left(\int_S |\nabla u|^2 + 2u^2 dA \right) \geq 0, \end{aligned}$$

where the equality in the second row follows from the Stoke's Theorem. The last equality holds if and only if $u = 0$. That is, D is positive. For the case that f is complex-valued, one can show it by arguing the real and imaginary parts. \square

For the Green function of the operator $-2(\Delta - 2)^{-1}$, we have

Proposition 2.2. *Let D be the operator above. Then there exists a Green function $G(w, z)$ for D satisfying:*

- (1) $G(w, z)$ is positive.
- (2) $G(w, z)$ is symmetric, i.e., $G(w, z) = G(z, w)$.

Proof. One can refer to [20] and [27]. \square

The Riemannian tensor of the Weil-Petersson metric. The curvature tensor is given by the following. Let μ_α, μ_β be two elements in the tangent space at X , and

$$g_{\alpha\bar{\beta}} = \int_X \mu_\alpha \cdot \overline{\mu_\beta} dA,$$

where dA is the area element for X .

Let us study the curvature tensor in these local coordinates. First of all, for the inverse of $(g_{i\bar{j}})$, we use the convention

$$g^{i\bar{j}} g_{k\bar{j}} = \delta_{ik}.$$

The curvature tensor is given by

$$R_{i\bar{j}k\bar{l}} = \frac{\partial^2 g_{i\bar{j}}}{\partial t^k \partial \bar{t}^l} - g^{s\bar{t}} \frac{\partial g_{i\bar{t}}}{\partial t^k} \frac{\partial g_{s\bar{j}}}{\partial \bar{t}^l}.$$

Since Ahlfors showed that the first derivatives of the metric tensor vanish at the base point X in these coordinates, at X we have

$$(1) \quad R_{i\bar{j}k\bar{l}} = \frac{\partial^2 g_{i\bar{j}}}{\partial t^k \partial \bar{t}^l}.$$

By the same argument in Kähler geometry we have

Proposition 2.3. *For any indices i, j, k, l , we have*

$$(1) \quad R_{i\bar{j}k\bar{l}} = R_{i\bar{j}l\bar{k}} = 0.$$

$$(2) \quad R_{i\bar{j}k\bar{l}} = -R_{i\bar{j}l\bar{k}}.$$

$$(3) \quad R_{i\bar{j}k\bar{l}} = R_{k\bar{j}i\bar{l}}.$$

$$(4) \quad R_{i\bar{j}k\bar{l}} = R_{i\bar{l}k\bar{j}}.$$

Proof. These follow from formula (1) and the first Bianchi identity (one can refer to [12]). \square

Now let us state Wolpert's curvature formula, which is crucial in the proof of theorem 1.1.

Theorem 2.4. (see [27]) *The curvature tensor satisfies*

$$R_{i\bar{j}k\bar{l}} = \int_X D(\mu_i \mu_{\bar{j}}) \cdot (\mu_k \mu_{\bar{l}}) dA + \int_X D(\mu_i \mu_{\bar{l}}) \cdot (\mu_k \mu_{\bar{j}}) dA.$$

Definition 2.5. Let μ_* be elements $\in T_X \text{Teich}(\mathcal{S})$. Set

$$(i\bar{j}, k\bar{l}) := \int_X D(\mu_i \mu_{\bar{j}}) \cdot (\mu_k \mu_{\bar{l}}) dA.$$

We close this section by rewriting theorem 2.4 as follows.

Theorem 2.6.

$$R_{i\bar{j}k\bar{l}} = (i\bar{j}, k\bar{l}) + (i\bar{l}, k\bar{j}).$$

3. CURVATURE OPERATOR ON SUBSPACES OF $\wedge^2 T_X^2 \text{Teich}(\mathcal{S})$

Before we study the curvature operator of $\text{Teich}(\mathcal{S})$, let us set some necessary notations. Let U be a neighborhood of X and $(t_1, t_2, \dots, t_{3g-3})$ be a local holomorphic coordinate on U , where $t_i = x_i + iy_i$ ($1 \leq i \leq 3g-3$). Then $(x_1, x_2, \dots, x_{3g-3}, y_1, y_2, \dots, y_{3g-3})$ is a real smooth coordinate in U . Furthermore, we have

$$\frac{\partial}{\partial x_i} = \frac{\partial}{\partial t_i} + \frac{\partial}{\partial \bar{t}_i}, \quad \frac{\partial}{\partial y_i} = i \left(\frac{\partial}{\partial t_i} - \frac{\partial}{\partial \bar{t}_i} \right).$$

Let $T \text{Teich}(\mathcal{S})$ be the real tangent bundle of $\text{Teich}(\mathcal{S})$ and $\wedge^2 T \text{Teich}(\mathcal{S})$ be the exterior wedge product of $T \text{Teich}(\mathcal{S})$ and itself. For any $X \in U$, we have

$$T_X \text{Teich}(\mathcal{S}) = \text{Span} \left\{ \frac{\partial}{\partial x_i}(X), \frac{\partial}{\partial y_j}(X) \right\}_{1 \leq i, j \leq 3g-3}$$

and

$$\wedge^2 T \text{Teich}(\mathcal{S}) = \text{Span} \left\{ \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}, \frac{\partial}{\partial x_k} \wedge \frac{\partial}{\partial y_l}, \frac{\partial}{\partial y_m} \wedge \frac{\partial}{\partial y_n} \right\}.$$

Set

$$\wedge^2 T_X^1 \text{Teich}(\mathcal{S}) := \text{Span} \left\{ \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} \right\},$$

$$\wedge^2 T_X^2 \text{Teich}(\mathcal{S}) := \text{Span} \left\{ \frac{\partial}{\partial x_k} \wedge \frac{\partial}{\partial y_l} \right\},$$

$$\wedge^2 T_X^3 \text{Teich}(\mathcal{S}) := \text{Span} \left\{ \frac{\partial}{\partial y_m} \wedge \frac{\partial}{\partial y_n} \right\}.$$

Hence,

$$\wedge^2 T_X \text{Teich}(\mathcal{S}) = \text{Span} \{ \wedge^2 T_X^1 \text{Teich}(\mathcal{S}), \wedge^2 T_X^2 \text{Teich}(\mathcal{S}), \wedge^2 T_X^3 \text{Teich}(\mathcal{S}) \}.$$

3.1. The curvature operator on $\wedge^2 T_X^1 \text{Teich}(\mathbb{S})$. Let $\sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}$ be an element in $\wedge^2 T_X^1 \text{Teich}(\mathbb{S})$, where a_{ij} are real. Set

$$F(z, w) = \sum_{i,j=1}^{3g-3} a_{ij} \mu_i(w) \cdot \overline{\mu_j(z)}.$$

The following proposition is influenced by theorem 4.1 in [15].

Proposition 3.1. *Let Q be the curvature operator and $D = -2(\Delta - 2)^{-1}$, where Δ is the Beltrami-Laplace operator on X . G is the Green function of D , and $\sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}$ is an element in $\wedge^2 T_X^1 \text{Teich}(\mathbb{S})$, where a_{ij} are real. Then we have*

$$\begin{aligned} & Q\left(\sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}, \sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}\right) \\ &= \int_X D(F(z, z) - \overline{F(z, z)})(F(z, z) - \overline{F(z, z)}) dA(z) \\ &- 2 \cdot \int_{X \times X} G(z, w) |F(z, w)|^2 dA(w) dA(z) \\ &+ 2 \cdot \Re \left\{ \int_{X \times X} G(z, w) F(z, w) F(w, z) dA(w) dA(z) \right\}, \end{aligned}$$

where $F(z, w) = \sum_{i,j=1}^{3g-3} a_{ij} \mu_i(w) \cdot \overline{\mu_j(z)}$.

Proof. Since $\frac{\partial}{\partial x_i} = \frac{\partial}{\partial t_i} + \frac{\partial}{\partial \bar{t}_i}$, from proposition 2.3,

$$\begin{aligned} & Q\left(\sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}, \sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}\right) \\ &= \sum_{i,j,k,l} a_{ij} a_{kl} (R_{i\bar{j}k\bar{l}} + R_{i\bar{j}l\bar{k}} + R_{i\bar{j}k\bar{l}} + R_{i\bar{j}l\bar{k}}) \\ &= \sum_{i,j,k,l} a_{ij} a_{kl} (R_{i\bar{j}k\bar{l}} - R_{i\bar{j}l\bar{k}} - R_{j\bar{i}k\bar{l}} + R_{j\bar{i}l\bar{k}}) \\ &= \sum_{i,j,k,l} a_{ij} a_{kl} ((i\bar{j}, k\bar{l}) + (i\bar{l}, k\bar{j}) - (i\bar{j}, l\bar{k}) - (i\bar{k}, l\bar{j}) \\ &- (j\bar{i}, k\bar{l}) - (j\bar{l}, k\bar{i}) + (j\bar{i}, l\bar{k}) + (j\bar{k}, l\bar{i})) \quad (\text{by theorem 2.6}) \\ &= \sum_{i,j,k,l} a_{ij} a_{kl} (i\bar{j} - j\bar{i}, k\bar{l} - l\bar{k}) \\ &+ \sum_{i,j,k,l} a_{ij} a_{kl} ((i\bar{l}, k\bar{j}) + (l\bar{i}, j\bar{k})) \\ &- \sum_{i,j,k,l} a_{ij} a_{kl} ((i\bar{k}, l\bar{j}) + (j\bar{l}, k\bar{i})). \end{aligned}$$

For the first term, from definition 2.5,

$$\begin{aligned}
& \sum_{i,j,k,l} a_{ij} a_{kl} (i\bar{j} - j\bar{i}, k\bar{l} - l\bar{k}) \\
&= \int_X D(\sum_{ij} a_{ij} \mu_i \bar{\mu}_j - \sum_{ij} a_{ij} \mu_j \bar{\mu}_i) (\sum_{ij} a_{ij} \mu_i \bar{\mu}_j - \sum_{ij} a_{ij} \mu_j \bar{\mu}_i) dA(z) \\
&= \int_X D(F(z, z) - \overline{F(z, z)}) (F(z, z) - \overline{F(z, z)}) dA(z).
\end{aligned}$$

For the second term, after applying the Green function G we have

$$\begin{aligned}
& \sum_{i,l} a_{ij} a_{kl} ((i\bar{l}, k\bar{j}) + (l\bar{i}, j\bar{k})) = 2 \cdot \Re \{ \sum_{i,l} a_{ij} a_{kl} ((i\bar{l}, k\bar{j})) \} \\
&= 2 \cdot \Re \{ \int_X D(\sum_i a_{ij} \mu_i \bar{\mu}_l) (\sum_k a_{kl} \mu_k \bar{\mu}_j) dA(z) \} \\
&= 2 \cdot \Re \{ \int_X \int_X G(w, z) \sum_i a_{ij} \mu_i(w) \overline{\mu_l(w)} (\sum_k a_{kl} \mu_k(z) \bar{\mu}_j(z)) dA(z) dA(w) \}.
\end{aligned}$$

From the definition of $F(z, w)$,

$$\begin{aligned}
& \sum_{i,j,k,l} a_{ij} a_{kl} ((i\bar{l}, k\bar{j}) + (l\bar{i}, j\bar{k})) \\
&= 2 \cdot \Re \{ \int_{X \times X} G(z, w) F(z, w) \overline{F(w, z)} dA(w) dA(z) \}.
\end{aligned}$$

For the last term, we use an argument similar to that for the second term,

$$\begin{aligned}
& \sum_{i,k} a_{ij} a_{kl} ((i\bar{k}, l\bar{j}) + (k\bar{i}, j\bar{l})) = 2 \cdot \Re \{ \sum_{i,k} a_{ij} a_{kl} ((i\bar{k}, l\bar{j})) \} \\
&= 2 \cdot \Re \{ \int_X D(\sum_i a_{ij} \mu_i \sum_k \overline{a_{kl} \mu_k}) (\mu_l \bar{\mu}_j) dA(z) \} \\
&= 2 \cdot \Re \{ \int_X \int_X G(w, z) \sum_i a_{ij} \mu_i(w) \overline{\sum_k a_{kl} \mu_k(w)} (\mu_l(z) \bar{\mu}_j(z)) dA(z) dA(w) \}
\end{aligned}$$

From the definition of $F(z, w)$,

$$\begin{aligned}
& \sum_{i,j,k,l} a_{ij} a_{kl} ((i\bar{k}, l\bar{j}) + (k\bar{i}, j\bar{l})) \\
&= 2 \cdot \Re \{ \int_{X \times X} G(z, w) F(z, w) \overline{F(z, w)} dA(w) dA(z) \} \\
&= 2 \cdot \int_{X \times X} G(z, w) |F(z, w)|^2 dA(w) dA(z).
\end{aligned}$$

The conclusion follows from combining the three terms above. \square

Using the Green function's positivity and symmetry,

Theorem 3.2. *Under the same conditions in proposition 3.1, Q is negative definite on $\wedge^2 T_X^1 \text{Teich}(\mathbb{S})$.*

Proof. By proposition 3.1 we have

$$\begin{aligned}
& Q\left(\sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}, \sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}\right) \\
&= \int_X D(F(z, z) - \overline{F(z, z)})(F(z, z) - \overline{F(z, z)}) dA(z) \\
&- 2 \cdot \left(\int_{X \times X} G(z, w) |F(z, w)|^2 dA(w) dA(z) \right) \\
&- \Re \left\{ \int_{X \times X} G(z, w) F(z, w) \overline{F(w, z)} dA(w) dA(z) \right\}.
\end{aligned}$$

For the first term, since $F(z, z) - \overline{F(z, z)} = 2i\Im\{F(z, z)\}$, by the positivity of the operator D ,

$$\begin{aligned}
& \int_X D(F(z, z) - \overline{F(z, z)})(F(z, z) - \overline{F(z, z)}) dA(z) \\
&= -4 \cdot \int_X D(\Im\{F(z, z)\})(\Im\{F(z, z)\}) dA(z) \leq 0.
\end{aligned}$$

For the second term, using the Cauchy-Schwarz inequality,

$$\begin{aligned}
& \left| \int_{X \times X} G(z, w) F(z, w) \overline{F(w, z)} dA(w) dA(z) \right| \\
&\leq \int_{X \times X} |G(z, w) F(z, w) \overline{F(w, z)}| dA(w) dA(z) \\
&\leq \sqrt{\int_{X \times X} |G(z, w)| |F(z, w)|^2 dA(w) dA(z)} \\
&\times \sqrt{\int_{X \times X} |G(z, w)| |F(w, z)|^2 dA(w) dA(z)} \\
&= \int_{X \times X} G(z, w) |F(z, w)|^2 dA(w) dA(z),
\end{aligned}$$

since G is positive and symmetric (see proposition 2.2).

Combining these three terms, we get that Q is non-positive definite on $\wedge^2 T_X^1 \text{Teich}(\mathbb{S})$.

Furthermore, equality holds precisely when

$$Q\left(\sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}, \sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}\right) = 0;$$

that is, there exists a constant complex number k such that both of the following hold:

$$\begin{cases} F(z, z) = \overline{F(z, z)}, \\ F(z, w) = k \cdot \overline{F(w, z)}. \end{cases}$$

If we let $z = w$, we get $k = 1$. Hence, the last equation is equivalent to

$$\sum_{ij} (a_{ij} - a_{ji}) \mu_i(w) \overline{\mu_j}(z) = 0.$$

Since $\{\mu_i\}_{i \geq 1}$ is a basis,

$$a_{ij} = a_{ji}.$$

This means $\sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} = 0$. That is, Q is negative definite on $\wedge^2 T_X^1 \text{Teich}(\mathcal{S})$. \square

3.2. The curvature operator on $\wedge^2 T_X^2 \text{Teich}(\mathcal{S})$. Let b_{ij} be real and $\sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j} \in \wedge^2 T_X^2 \text{Teich}(\mathcal{S})$. Set

$$H(z, w) = \sum_{i,j=1}^{3g-3} b_{ij} \mu_i(w) \cdot \overline{\mu_j(z)}.$$

Using a similar computation in proposition 3.1, the formula for the curvature operator on $\wedge^2 T_X^2 \text{Teich}(\mathcal{S})$ is given as follows.

Proposition 3.3. *Let Q be the curvature operator and D be the same operator as shown in proposition 3.1. Let $\sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}$ be an element in $\wedge^2 T_X^2 \text{Teich}(\mathcal{S})$, where b_{ij} are real. Then we have*

$$\begin{aligned} & Q\left(\sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}, \sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}\right) \\ &= - \int_X D(H(z, z) + \overline{H(z, z)})(H(z, z) + \overline{H(z, z)}) dA(z) \\ &\quad - 2 \cdot \int_{X \times X} G(z, w) |H(z, w)|^2 dA(w) dA(z) \\ &\quad - 2 \cdot \Re \left\{ \int_{X \times X} G(z, w) H(z, w) \overline{H(w, z)} dA(w) dA(z), \right\} \end{aligned}$$

where $H(z, w) = \sum_{i,j=1}^{3g-3} b_{ij} \mu_i(w) \cdot \overline{\mu_j(z)}$.

Proof. Since $\frac{\partial}{\partial x_i} = \frac{\partial}{\partial t_i} + \frac{\partial}{\partial \bar{t}_i}$ and $\frac{\partial}{\partial y_i} = \mathbf{i}(\frac{\partial}{\partial t_i} - \frac{\partial}{\partial \bar{t}_i})$, from proposition 2.3,

$$\begin{aligned} & Q\left(\sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}, \sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}\right) \\ &= - \sum_{i,j,k,l} b_{ij} b_{kl} (R_{i\bar{j}k\bar{l}} - R_{i\bar{j}l\bar{k}} - R_{j\bar{i}k\bar{l}} + R_{j\bar{i}l\bar{k}}) \\ &= - \sum_{i,j,k,l} b_{ij} b_{kl} (R_{i\bar{j}k\bar{l}} + R_{i\bar{j}l\bar{k}} + R_{j\bar{i}k\bar{l}} + R_{j\bar{i}l\bar{k}}). \\ &= - \sum_{i,j,k,l} b_{ij} b_{kl} ((i\bar{j}, k\bar{l}) + (i\bar{l}, k\bar{j}) + (j\bar{i}, l\bar{k}) + (j\bar{k}, l\bar{i})) \quad (\text{by theorem 2.6}) \\ &\quad + (j\bar{l}, k\bar{i}) + (j\bar{i}, k\bar{l}) + (j\bar{k}, l\bar{i}) + (j\bar{l}, l\bar{i})) \\ &= - \sum_{i,j,k,l} b_{ij} b_{kl} (i\bar{j} + j\bar{i}, k\bar{l} + l\bar{k}) \\ &\quad - \sum_{i,j,k,l} b_{ij} b_{kl} ((i\bar{l}, k\bar{j}) + (l\bar{i}, j\bar{k})) \\ &\quad - \sum_{i,j,k,l} b_{ij} b_{kl} ((i\bar{k}, l\bar{j}) + (j\bar{l}, k\bar{i})). \end{aligned}$$

For the first term, from definition 2.5,

$$\begin{aligned}
& - \sum_{i,j,k,l} b_{ij} b_{kl} (i\bar{j} + j\bar{i}, k\bar{l} + l\bar{k}) \\
& = - \int_X D \left(\sum_{ij} b_{ij} \mu_i \overline{\mu_j} + \sum_{ij} b_{ij} \mu_j \overline{\mu_i} \right) \left(\sum_{ij} b_{ij} \mu_i \overline{\mu_j} + \sum_{ij} b_{ij} \mu_j \overline{\mu_i} \right) dA(z) \\
& = - \int_X D(H(z, z) + \overline{H(z, z)})(H(z, z) + \overline{H(z, z)}) dA(z).
\end{aligned}$$

For the second term and the third term, using the same argument in the proof of proposition 3.1, we have

$$\begin{aligned}
& \sum_{i,j,k,l} b_{ij} b_{kl} ((i\bar{l}, k\bar{j}) + (l\bar{i}, j\bar{k})) \\
& = 2 \cdot \Re \left\{ \int_{X \times X} G(z, w) H(z, w) H(w, z) dA(w) dA(z) \right\}
\end{aligned}$$

and

$$\begin{aligned}
& \sum_{i,j,k,l} b_{ij} b_{kl} ((i\bar{k}, l\bar{j}) + (k\bar{i}, j\bar{l})) \\
& = 2 \cdot \int_{X \times X} G(z, w) |H(z, w)|^2 dA(w) dA(z).
\end{aligned}$$

Combining these three terms we get the proposition. \square

Using the same method in theorem 3.2, one can prove the following non-positivity result.

Theorem 3.4. *Under the same conditions of proposition 3.3, then Q is non-positive definite on $\wedge^2 T_X^2 \text{Teich}(S)$, and the zero level subsets of $Q(\cdot, \cdot)$ are $\{\sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}; b_{ij} = -b_{ji}\}$.*

Proof. Let $\sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}$ be an element in $\wedge^2 T_X^2 \text{Teich}(S)$. From proposition 3.3 we have

$$\begin{aligned}
& Q \left(\sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}, \sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j} \right) \\
& = - \int_X D(H(z, z) + \overline{H(z, z)})(H(z, z) + \overline{H(z, z)}) dA(z) \\
& \quad - 2 \left(\int_{X \times X} G(z, w) |H(z, w)|^2 dA(w) dA(z) \right) \\
& \quad + \Re \left\{ \int_{X \times X} G(z, w) H(z, w) H(w, z) dA(w) dA(z) \right\}.
\end{aligned}$$

For the first term, since $H(z, z) + \overline{H(z, z)} = 2 \cdot \Re\{H(z, z)\}$, by the positivity of the operator D ,

$$\begin{aligned}
& - \int_X D(H(z, z) + \overline{H(z, z)})(H(z, z) + \overline{H(z, z)}) dA(z) \\
& = -4 \int_X D(\Re\{H(z, z)\})(\Re\{H(z, z)\}) dA(z) \leq 0.
\end{aligned}$$

For the second term, using the Cauchy-Schwarz inequality,

$$\begin{aligned}
& \left| \int_{X \times X} G(z, w) H(z, w) H(w, z) dA(w) dA(z) \right| \\
& \leq \int_{X \times X} |G(z, w) H(z, w) H(w, z)| dA(w) dA(z) \\
& \leq \sqrt{\int_{X \times X} |G(z, w)| |H(z, w)|^2 dA(w) dA(z)} \\
& \quad \times \sqrt{\int_{X \times X} |G(z, w)| |H(w, z)|^2 dA(w) dA(z)} \\
& = \int_{X \times X} G(z, w) |H(z, w)|^2 dA(w) dA(z),
\end{aligned}$$

since G is positive and symmetric.

Combining these two terms, we get Q is non-positive on $\wedge^2 T_X^2 \text{Teich}(\mathbb{S})$.

Using the same argument as in the proof of theorem 3.2,

$$Q\left(\sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}, \sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}\right) = 0$$

if and only if there exists a constant complex number k such that both of the following hold:

$$\begin{cases} H(z, z) = -\overline{H(z, z)}, \\ H(z, w) = k \cdot \overline{H(w, z)}. \end{cases}$$

If we let $z = w$, we get $k = -1$. Hence, the last equation is equivalent to

$$\sum_{ij} (b_{ij} + b_{ji}) \mu_i(w) \overline{\mu_j}(z) = 0.$$

Since $\{\mu_i\}_{i \geq 1}$ is a basis,

$$b_{ij} = -b_{ji}.$$

□

3.3. The curvature operator on $\wedge^2 T_X^3 \text{Teich}(\mathbb{S})$. Let \mathbf{J} be the almost complex structure on $\text{Teich}(\mathbb{S})$. Since $\{t_i\}$ is a holomorphic coordinate, we have

$$\mathbf{J} \frac{\partial}{\partial x_i} = \frac{\partial}{\partial y_i} \quad \mathbf{J} \frac{\partial}{\partial y_i} = -\frac{\partial}{\partial x_i}.$$

Since the Weil-Petersson metric is a Kähler metric, \mathbf{J} is an isometry on the tangent space. In particular we have

$$\begin{aligned}
R(V_1, V_2, V_3, V_4) &= R(\mathbf{J}V_1, \mathbf{J}V_2, \mathbf{J}V_3, \mathbf{J}V_4) \\
&= R(\mathbf{J}V_1, \mathbf{J}V_2, V_3, V_4) = R(V_1, V_2, \mathbf{J}V_3, \mathbf{J}V_4),
\end{aligned}$$

where R is the curvature tensor and V_i are real tangent vectors in $T_X \text{Teich}(\mathbb{S})$. Once can refer to [12] for more details.

Let $C = \sum_{ij} c_{ij} \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}$ be an element in $\wedge^2 T_X^3 \text{Teich}(\mathbb{S})$, where c_{ij} are real. Set

$$K(z, w) = \sum_{i,j=1}^{3g-3} c_{ij} \mu_i(w) \cdot \overline{\mu_j(z)}.$$

Proposition 3.5. *Let Q be the curvature operator, and $\sum_{ij} c_{ij} \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}$ be an element in $\wedge^2 T_X^3 \text{Teich}(\mathbb{S})$. Then we have*

$$\begin{aligned} & Q\left(\sum_{ij} c_{ij} \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}, \sum_{ij} c_{ij} \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}\right) \\ &= \int_X D(K(z, z) - \overline{K(z, z)})(K(z, z) - \overline{K(z, z)}) dA(z) \\ &- 2 \cdot \int_{X \times X} G(z, w) |K(z, w)|^2 dA(w) dA(z) \\ &+ 2 \cdot \Re \left\{ \int_{X \times X} G(z, w) K(z, w) K(w, z) dA(w) dA(z) \right\}. \end{aligned}$$

Proof. Since $\frac{\partial}{\partial y_i} = \mathbf{J} \frac{\partial}{\partial x_i} = \mathbf{J} \frac{\partial}{\partial t_i} + \mathbf{J} \frac{\partial}{\partial \bar{t}_i}$ and \mathbf{J} is an isometry, by proposition 2.3,

$$\begin{aligned} & Q\left(\sum_{ij} c_{ij} \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}, \sum_{ij} c_{ij} \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}\right) \\ &= \sum_{i,j,k,l} c_{ij} c_{kl} (R_{ij\bar{k}l} + R_{ij\bar{l}k} + R_{i\bar{j}kl} + R_{i\bar{j}lk}) \\ &= Q\left(\sum_{ij} c_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}, \sum_{ij} c_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}\right). \end{aligned}$$

By proposition 3.1,

$$\begin{aligned} & Q\left(\sum_{ij} c_{ij} \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}, \sum_{ij} c_{ij} \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}\right) \\ &= \int_X D(K(z, z) - \overline{K(z, z)})(K(z, z) - \overline{K(z, z)}) dA(z) \\ &- 2 \cdot \int_{X \times X} G(z, w) |K(z, w)|^2 dA(w) dA(z) \\ &+ 2 \cdot \Re \left\{ \int_{X \times X} G(z, w) K(z, w) K(w, z) dA(w) dA(z) \right\}. \end{aligned}$$

□

Using the same argument as in the proof of theorem 3.2 we have

Theorem 3.6. *Let Q be the curvature operator as above, then Q is a negative definite operator on $\wedge^2 T_X^3 \text{Teich}(\mathbb{S})$.*

4. CURVATURE OPERATOR ON $\wedge^2 T_X \text{Teich}(\mathbb{S})$

Every element in $\wedge^2 T_X \text{Teich}(\mathbb{S})$ can be represented by $\sum_{ij} (a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} + b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j} + c_{ij} \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j})$.

Proposition 4.1. *Let Q be the curvature operator. Then*

$$\begin{aligned} & Q\left(\sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} + b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j} + c_{ij} \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}, \right. \\ & \left. \sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} + b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j} + c_{ij} \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}\right) \\ &= Q\left(\sum_{ij} d_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} + b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}, \sum_{ij} d_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} + b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}\right), \end{aligned}$$

where $d_{ij} = a_{ij} + c_{ij}$.

Proof. Since the almost complex structure \mathbf{J} is an isometry and $\mathbf{J} \frac{\partial}{\partial x_i} = \frac{\partial}{\partial y_i}$, we have

$$\begin{aligned} Q\left(\frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}, \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}\right) &= R\left(\frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_j}, \mathbf{J} \frac{\partial}{\partial x_i}, \mathbf{J} \frac{\partial}{\partial x_j}\right) \\ &= R\left(\frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_j}, \frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_j}\right) \end{aligned}$$

and

$$\begin{aligned} Q\left(\frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}, \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}\right) &= R\left(\frac{\partial}{\partial x_i}, \frac{\partial}{\partial y_j}, \mathbf{J} \frac{\partial}{\partial x_i}, \mathbf{J} \frac{\partial}{\partial x_j}\right) \\ &= R\left(\frac{\partial}{\partial x_i}, \frac{\partial}{\partial y_j}, \frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_j}\right). \end{aligned}$$

The conclusion follows by expanding Q and applying the two equations above. \square

If one wants to determine whether the curvature operator Q is non-positive definite on $\wedge^2 T_X \text{Teich}(\mathbb{S})$, by proposition 4.1 it is sufficient to see if Q is non-positive definite on $\text{Span}\{\wedge^2 T_X^1 \text{Teich}(\mathbb{S}), \wedge^2 T_X^2 \text{Teich}(\mathbb{S})\}$.

Proposition 4.2. *Let Q be the curvature operator, $\sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}$ be an element in $\wedge^2 T_X^1 \text{Teich}(\mathbb{S})$, and $\sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}$ be an element in $\wedge^2 T_X^2 \text{Teich}(\mathbb{S})$. Then we have*

$$\begin{aligned} & Q\left(\sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}, \sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}\right) \\ &= \mathbf{i} \cdot \int_X D(F(z, z) - \overline{F(z, z)}) \cdot (H(z, z) + \overline{H(z, z)}) dA(z) \\ &\quad - 2 \cdot \Im \left\{ \int_{X \times X} G(z, w) F(z, w) \overline{H(z, w)} dA(w) dA(z) \right\} \\ &\quad - 2 \cdot \Im \left\{ \int_{X \times X} G(z, w) F(z, w) H(w, z) dA(w) dA(z) \right\}. \end{aligned}$$

Proof. Since $\frac{\partial}{\partial x_i} = \frac{\partial}{\partial t_i} + \frac{\partial}{\partial \bar{t}_i}$ and $\frac{\partial}{\partial y_i} = \mathbf{i}(\frac{\partial}{\partial t_i} - \frac{\partial}{\partial \bar{t}_i})$, by proposition 2.3,

$$\begin{aligned}
& Q\left(\sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}, \sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}\right) \\
&= (-\mathbf{i}) \sum_{i,j,k,l} a_{ij} b_{kl} (-R_{i\bar{j}k\bar{l}} + R_{i\bar{j}k\bar{l}} - R_{i\bar{j}k\bar{l}} + R_{i\bar{j}k\bar{l}}) \\
&= (-\mathbf{i}) \sum_{i,j,k,l} a_{ij} b_{kl} (-R_{i\bar{j}k\bar{l}} - R_{i\bar{j}l\bar{k}} + R_{j\bar{i}k\bar{l}} + R_{j\bar{i}l\bar{k}}) \\
&= (-\mathbf{i}) \sum_{i,j,k,l} a_{ij} b_{kl} (-(i\bar{j}, k\bar{l}) - (i\bar{l}, k\bar{j}) - (i\bar{j}, l\bar{k}) - (i\bar{k}, l\bar{j}) \\
&\quad + (j\bar{l}, k\bar{l}) + (j\bar{l}, k\bar{i}) + (j\bar{i}, l\bar{k}) + (j\bar{k}, l\bar{i})) \quad (\text{by theorem 2.6}) \\
&= (-\mathbf{i}) \sum_{i,j,k,l} a_{ij} b_{kl} (j\bar{i} - i\bar{j}, k\bar{l} + l\bar{k}) \\
&\quad + (-\mathbf{i}) \sum_{i,j,k,l} a_{ij} b_{kl} (-(i\bar{l}, k\bar{j}) + (l\bar{i}, j\bar{k})) \\
&\quad + (-\mathbf{i}) \sum_{i,j,k,l} a_{ij} b_{kl} (-(i\bar{k}, l\bar{j}) + (j\bar{l}, k\bar{i})).
\end{aligned}$$

For the first term, by definition 2.5,

$$\begin{aligned}
& \sum_{i,j,k,l} a_{ij} b_{kl} (j\bar{i} - i\bar{j}, k\bar{l} + l\bar{k}) \\
&= \int_X D\left(\sum_{ij} a_{ij} \bar{\mu}_i \mu_j - \sum_{ij} a_{ij} \mu_i \bar{\mu}_j\right) \left(\sum_{kl} b_{kl} \mu_k \bar{\mu}_l + \sum_{kl} b_{kl} \mu_l \bar{\mu}_k\right) dA(z) \\
&= \int_X D(\overline{F(z, z)} - F(z, z))(H(z, z) + \overline{H(z, z)}) dA(z).
\end{aligned}$$

For the second term, by using the Green function G of D ,

$$\begin{aligned}
& \sum_{i,l} a_{ij} b_{kl} (-(i\bar{l}, k\bar{j}) + (l\bar{i}, j\bar{k})) \\
&= 2\mathbf{i} \cdot \Im\left\{\sum_{i,l} a_{ij} b_{kl} (-(i\bar{l}, k\bar{j})\right\} \\
&= -2\mathbf{i} \cdot \Im\left\{\int_X D\left(\sum_i a_{ij} \mu_i \bar{\mu}_l\right) \left(\sum_k b_{kl} \mu_k \bar{\mu}_j\right) dA(z)\right\} \\
&= -2\mathbf{i} \cdot \Im\left\{\int_X \int_X G(w, z) \sum_i a_{ij} \mu_i(w) \overline{\mu_l(w)} \left(\sum_k b_{kl} \mu_k(z) \bar{\mu}_j(z)\right) dA(z) dA(w)\right\} \\
&= -2\mathbf{i} \cdot \Im\left\{\int_{X \times X} G(z, w) F(z, w) H(w, z) dA(w) dA(z)\right\}.
\end{aligned}$$

For the last term,

$$\begin{aligned}
& \sum_{i,k} a_{ij} b_{kl} (-(i\bar{k}, l\bar{j}) + (k\bar{i}, j\bar{l})) = -2\mathbf{i} \cdot \Im \left\{ \sum_{i,k} a_{ij} b_{kl} (i\bar{k}, l\bar{j}) \right\} \\
&= -2\mathbf{i} \cdot \Im \left\{ \int_X D \left(\sum_i a_{ij} \mu_i \overline{\sum_k b_{kl} \mu_k} (\mu_i \overline{\mu_j}) \right) dA(z) \right\} \\
&= -2\mathbf{i} \cdot \Im \left\{ \int_X \int_X G(w, z) \sum_i a_{ij} \mu_i(w) \overline{\sum_k b_{kl} \mu_k(w)} (\mu_i(z) \overline{\mu_j(z)}) dA(z) dA(w) \right\} \\
&= -2\mathbf{i} \cdot \Im \left\{ \int_{X \times X} G(z, w) F(z, w) \overline{H(z, w)} dA(w) dA(z) \right\}.
\end{aligned}$$

Combining these three terms above, we get the lemma. \square

The following proposition will give the formula for curvature operator Q on $Span\{\wedge^2 T_X^1 \text{Teich}(S), \wedge^2 T_X^2 \text{Teich}(S)\}$. Setting

$$A = \sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}, \quad B = \sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j},$$

on $Span\{\wedge^2 T_X^1 \text{Teich}(S), \wedge^2 T_X^2 \text{Teich}(S)\}$, we have

Proposition 4.3. *Let Q be the curvature operator on $\text{Teich}(S)$. Let $A = \sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}$ and $B = \sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}$. Then we have*

$$\begin{aligned}
& Q(A + B, A + B) = \\
& -4 \int_X D(\Im\{F(z, z) + \mathbf{i}H(z, z)\}) \cdot (\Im\{F(z, z) + \mathbf{i}H(z, z)\}) dA(z) \\
& -2 \int_{X \times X} G(z, w) |F(z, w) + \mathbf{i}H(z, w)|^2 dA(w) dA(z) \\
& +2 \Re \left\{ \int_{X \times X} G(z, w) (F(z, w) + \mathbf{i}H(z, w)) (F(w, z) + \mathbf{i}H(w, z)) dA(w) dA(z) \right\},
\end{aligned}$$

where $F(z, w) = \sum_{i,j=1}^{3g-3} a_{ij} \mu_i(w) \cdot \overline{\mu_j(z)}$ and $H(z, w) = \sum_{i,j=1}^{3g-3} b_{ij} \mu_i(w) \cdot \overline{\mu_j(z)}$.

Proof. Since $Q(A, B) = Q(B, A)$,

$$Q(A + B, A + B) = Q(A, A) + 2Q(A, B) + Q(B, B).$$

By proposition 3.1, proposition 3.3, and proposition 4.2 we have

$$\begin{aligned}
& Q(A + B, A + B) \\
&= \left(\int_X D(F(z, z) - \overline{F(z, z)})(F(z, z) - \overline{F(z, z)})dA(z) \right. \\
&\quad - \int_X D(H(z, z) + \overline{H(z, z)})(H(z, z) + \overline{H(z, z)})dA(z) \\
&\quad + 2\mathbf{i} \cdot \int_X D(F(z, z) - \overline{F(z, z)})(H(z, z) + \overline{H(z, z)})dA(z) \Big) \\
&\quad - 2 \cdot \int_{X \times X} G(z, w) |F(z, w)|^2 dA(w) dA(z) \\
&\quad - 2 \cdot \int_{X \times X} G(z, w) |H(z, w)|^2 dA(w) dA(z) \\
&\quad - 4 \cdot \Im \left\{ \int_{X \times X} G(z, w) F(z, w) \overline{H(z, w)} dA(w) dA(z) \right\} \\
&\quad + 2 \cdot \Re \left\{ \int_{X \times X} G(z, w) F(z, w) F(w, z) dA(w) dA(z) \right\} \\
&\quad - 2 \cdot \Re \left\{ \int_{X \times X} G(z, w) H(z, w) H(w, z) dA(w) dA(z) \right\} \\
&\quad - 4 \cdot \Im \left\{ \int_{X \times X} G(z, w) F(z, w) H(w, z) dA(w) dA(z) \right\}.
\end{aligned}$$

The sum of the first three terms is exactly

$$-4 \int_X D(\Im\{F(z, z) + \mathbf{i}H(z, z)\}) \cdot (\Im\{F(z, z) + \mathbf{i}H(z, z)\})dA(z).$$

Just as $|a + \mathbf{i}b|^2 = |a|^2 + |b|^2 + 2 \cdot \Im(a \cdot \overline{b})$, where a and b are two complex numbers, the sum of the second three terms is exactly

$$-2 \cdot \int_{X \times X} G(z, w) |F(z, w) + \mathbf{i}H(z, w)|^2 dA(w) dA(z).$$

For the last three terms, since

$$\Im(F(z, w) \cdot H(w, z)) = -\Re(F(z, w) \cdot (\mathbf{i}H(w, z))),$$

the sum is exactly

$$2 \cdot \Re \left\{ \int_{X \times X} G(z, w) (F(z, w) + \mathbf{i}H(z, w))(F(w, z) + \mathbf{i}H(w, z)) dA(w) dA(z) \right\}.$$

□

Furthermore, we have

Theorem 4.4. *Under the same conditions as in proposition 4.3, Q is non-positive definite on $\text{Span}\{\wedge^2 T_X^1 \text{Teich}(\mathbf{S}), \wedge^2 T_X^2 \text{Teich}(\mathbf{S})\}$, and the zero level subsets of $Q(\cdot, \cdot)$ are $\{\sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}; b_{ij} = -b_{ji}\}$ in $\text{Span}\{\wedge^2 T_X^1 \text{Teich}(\mathbf{S}), \wedge^2 T_X^2 \text{Teich}(\mathbf{S})\}$.*

Proof. Let us estimate the terms in proposition 4.3 separately. For the first term, since D is a positive operator,

$$-\int_X D(\Im\{F(z, z) + \mathbf{i}H(z, z)\}) \cdot (\Im\{F(z, z) + \mathbf{i}H(z, z)\}) dA(z) \leq 0.$$

For the third term, by the Cauchy-Schwarz inequality,

$$\begin{aligned} & \left| \int_{X \times X} G(z, w)(F(z, w) + \mathbf{i}H(z, w))(F(w, z) + \mathbf{i}H(w, z)) dA(w) dA(z) \right| \\ & \leq \int_{X \times X} G(z, w) |(F(z, w) + \mathbf{i}H(z, w))(F(w, z) + \mathbf{i}H(w, z))| dA(w) dA(z) \\ & \leq \sqrt{\int_{X \times X} G(z, w) |(F(z, w) + \mathbf{i}H(z, w))|^2 dA(w) dA(z)} \\ & \quad \times \sqrt{\int_{X \times X} G(z, w) |(F(w, z) + \mathbf{i}H(w, z))|^2 dA(w) dA(z)} \\ & = \int_{X \times X} G(z, w) |(F(z, w) + \mathbf{i}H(z, w))|^2 dA(w) dA(z). \end{aligned}$$

The last equality follows from $G(z, w) = G(w, z)$.

Combining the two inequalities above and the second term in proposition 4.3, we see that on $\text{Span}\{\wedge^2 T_X^1 \text{Teich}(S), \wedge^2 T_X^2 \text{Teich}(S)\}$ Q is non-positive definite. Furthermore, $Q(A + B, A + B) = 0$ if and only if there exists a constant k such that both of the following hold:

$$\begin{cases} \text{Im}\{F(z, z) + \mathbf{i}H(z, z)\} = 0, \\ F(z, w) + \mathbf{i}H(z, w) = k \cdot \overline{(F(w, z) + \mathbf{i}H(w, z))}. \end{cases}$$

If we let $z = w$, we get $k = 1$. Hence, the second equation above is

$$\sum_{ij} (a_{ij} - a_{ji} + \mathbf{i}(b_{ij} + b_{ji})) \mu_i(w) \overline{\mu_j(z)} = 0.$$

Since $\{\mu_i\}_{i \geq 1}$ is a basis,

$$a_{ij} = a_{ji}, \quad b_{ij} = -b_{ji}.$$

That is, $A = 0$ and $B = \sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}$, where $b_{ij} = -b_{ji}$.

Conversely, if $A = 0$ and $B \in \{\sum_{ij} b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}; b_{ij} = -b_{ji}\}$, it is not hard to apply proposition 4.3 to show that $Q(A + B, A + B) = 0$. \square

Before we prove the main theorem, let us define a natural action of \mathbf{J} on $\wedge^2 T_X \text{Teich}(S)$ by

$$\begin{cases} \mathbf{J} \circ \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} := \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}, \\ \mathbf{J} \circ \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j} := -\frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial x_j} = \frac{\partial}{\partial x_j} \wedge \frac{\partial}{\partial y_i}, \\ \mathbf{J} \circ \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j} := \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}, \end{cases}$$

and extend it linearly. It is easy to see that $\mathbf{J} \circ \mathbf{J} = id$.

Now we are ready to prove theorem 1.1.

Proof of Theorem 1.1. It follows from proposition 4.1 and theorem 4.4 that Q is non-positive definite.

If $A = C - \mathbf{J} \circ C$ for some $C \in \wedge^2 T_X \text{Teich}(S)$. Then it is easy to see that $Q(A, A) = 0$ since \mathbf{J} is an isometry.

Assume that $A \in \wedge^2 T_X \text{Teich}(\mathbb{S})$ such that $Q(A, A) = 0$. Since $\wedge^2 T \text{Teich}(\mathbb{S}) = \text{Span}\{\frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}, \frac{\partial}{\partial x_k} \wedge \frac{\partial}{\partial y_l}, \frac{\partial}{\partial y_m} \wedge \frac{\partial}{\partial y_n}\}$, there exists a_{ij}, b_{ij} , and c_{ij} such that

$$A = \sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} + b_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j} + c_{ij} \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}.$$

Since $Q(A, A) = 0$, by proposition 4.1 and theorem 4.4 we must have

$$a_{ij} + c_{ij} = a_{ji} + c_{ji}, b_{ij} = -b_{ji}.$$

That is,

$$a_{ij} - a_{ji} = -(c_{ij} - c_{ji}), b_{ij} = -b_{ji}.$$

Set

$$C = \sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} + \frac{b_{ij}}{2} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}.$$

Claim: $A = C - \mathbf{J} \circ C$.

Since $\sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} = \sum_{i < j} (a_{ij} - a_{ji}) \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}$, we have

$$\begin{aligned} \mathbf{J} \circ \sum_{ij} a_{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j} &= \sum_{i < j} (a_{ij} - a_{ji}) \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j} \\ &= - \sum_{i < j} (c_{ij} - c_{ji}) \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j} = - \sum c_{ij} \frac{\partial}{\partial y_i} \wedge \frac{\partial}{\partial y_j}. \end{aligned}$$

Similarly,

$$\mathbf{J} \circ \sum \left(\frac{b_{ij}}{2} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j} \right) = - \sum \frac{b_{ij}}{2} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_j}.$$

The claim follows from the two equations above. \square

5. HARMONIC MAPS INTO $\text{Teich}(\mathbb{S})$

In this section we study the twist-harmonic maps from some domains into the Teichmüller space. Before we go to the rank-one hyperbolic space case, let us state the following lemma, which is influenced by lemma 5 in [32].

Lemma 5.1. *Let M be either $H_{Q,m} = Sp(m, 1)/Sp(m) \cdot Sp(1)$ or $H_{O,2} = F_4^{-20}/SO(9)$. Then the rank-one Hyperbolic space M cannot be totally geodesically immersed into $\text{Teich}(\mathbb{S})$.*

Proof. For $H_{Q,m} = Sp(m, 1)/Sp(m)$, we assume that there is a totally geodesic immersion of $H_{Q,m}$ into $\text{Teich}(\mathbb{S})$. We may select $p \in H_{Q,m}$. Choose a quaternionic line l_Q on $T_p H_{Q,m}$, and we may assume that l_Q is spanned over \mathbb{R} by v, Iv, Jv , and Kv . Without loss of generality, we may assume that J on $l_Q \subset T_p H_{Q,m}$ is the same as the complex structure on $\text{Teich}(\mathbb{S})$. Choose an element

$$v \wedge Jv + Kv \wedge Iv \in \wedge^2 T_p H_{Q,m}.$$

Let $Q^{H_{Q,m}}$ be the curvature operator on $H_{Q,m}$.

$$\begin{aligned} Q^{H_{Q,m}}(v \wedge Jv + Kv \wedge Iv, v \wedge Jv + Kv \wedge Iv) &= \\ R^{H_{Q,m}}(v, Jv, v, Jv) + R^{H_{Q,m}}(Kv, Iv, Kv, Iv) + 2 \cdot R^{H_{Q,m}}(v, Jv, Kv, Iv). \end{aligned}$$

Since I is an isometry, we have

$$\begin{aligned} R^{H_{Q,m}}(Kv, Iv, Kv, Iv) &= R^{H_{Q,m}}(IKv, IIv, IKv, IIv) \\ &= R^{H_{Q,m}}(-Jv, -v, -Jv, -v) \\ &= R^{H_{Q,m}}(v, Jv, v, Jv). \end{aligned}$$

Similarly,

$$\begin{aligned} R^{H_{Q,m}}(v, Jv, Kv, Iv) &= R^{H_{Q,m}}(v, Jv, IKv, IIv) \\ &= R^{H_{Q,m}}(v, Jv, -Jv, -v) \\ &= -R^{H_{Q,m}}(v, Jv, v, Jv). \end{aligned}$$

Combining the terms above, we have

$$Q^{H_{Q,m}}(v \wedge Jv + Kv \wedge Iv, v \wedge Jv + Kv \wedge Iv) = 0.$$

Since f is a geodesical immersion,

$$Q^{\text{Teich}(S)}(v \wedge Jv + Kv \wedge Iv, v \wedge Jv + Kv \wedge Iv) = 0.$$

On the other hand, by theorem 1.1, there exists C such that

$$v \wedge Jv + Kv \wedge Iv = C - \mathbf{J} \circ C.$$

Hence,

$$\begin{aligned} (2) \quad & \mathbf{J} \circ (v \wedge Jv + Kv \wedge Iv) \\ &= \mathbf{J} \circ (C - \mathbf{J} \circ C) = \mathbf{J} \circ C - \mathbf{J} \circ \mathbf{J} \circ C = \mathbf{J} \circ C - C \\ &= -(v \wedge Jv + Kv \wedge Iv). \end{aligned}$$

On the other hand, since \mathbf{J} is the same as J in $H_{Q,m}$, we also have

$$\begin{aligned} (3) \quad & \mathbf{J} \circ (v \wedge Jv + Kv \wedge Iv) = (Jv \wedge JJv + JKv \wedge JIv) \\ &= Jv \wedge (-v) + Iv \wedge (-Kv) = v \wedge Jv + Kv \wedge Iv. \end{aligned}$$

From equations (2) and (3) we get

$$v \wedge Jv + Kv \wedge Iv = 0,$$

which is a contradiction since l_Q is spanned over R by v, Iv, Jv , and Kv .

In the case of the Cayley hyperbolic plane $H_{O,2} = F_4^{20}/SO(9)$, the argument is similar by replacing a quaternionic line by a Cayley line ([4]). \square

Now we are ready to prove theorem 1.2.

Proof of theorem 1.2. Since the sectional curvature operator on $\text{Teich}(S)$ is non-positive definite, $\text{Teich}(S)$ also has non-positive Riemannian sectional curvature in the complexified sense as stated in [19]. Suppose that f is not constant. From theorem 2 in [19] (also see [6]), we know that f should be a totally geodesic immersion, which contradicts lemma 5.1. Hence, f must be a constant. \square

Remark 5.1. In [32] it is shown that the image of any homomorphism ρ from Γ to $\text{Mod}(S)$ is finite. Hence, $\rho(\Gamma)$ must have a fixed point in $\text{Teich}(S)$ from the Nielsen realization theorem (one can see [14, 30]). If we assume that there exists a twist harmonic map f with respect to this homomorphism, then by theorem 1.2 we know $\rho(\Gamma) \subset \text{Mod}(S)$ will fix the point $f(G/\Gamma) \in \text{Teich}(S)$.

Remark 5.2. Conversely, if one can prove that for any homomorphism ρ from Γ to $\text{Mod}(S)$ there exists a twist harmonic map f from G into the completion $\overline{\text{Teich}(S)}$ of $\text{Teich}(S)$ such that the image $f(G) \subset \overline{\text{Teich}(S)}$, theorem 1.2 tells us that the image $\rho(\Gamma)$ fixes a point in $\text{Teich}(S)$; hence, the image $\rho(\Gamma)$ is finite because $\text{Mod}(S)$ acts properly on $\text{Teich}(S)$.

REFERENCES

- [1] Lars V. Ahlfors, *Some remarks on Teichmüller space of Riemann surfaces*, Ann. of Math. (2) **74** (1961), 171–191, MR 0204641, Zbl 0146.30602.
- [2] C. S. Aravinda and F. T. Farrell, *Nonpositivity: curvature vs. curvature operator*, Proc. Amer. Math. Soc. **133** (2005), no. 1, 191–192, MR 2085169, Zbl 1065.53031.
- [3] J. Brock and B. Farb, *Curvature and rank of Teichmüller space*, Amer. J. Math. **128** (2006), no. 1, 1–22, MR 2197066, Zbl 1092.32008.
- [4] I. Chavel, *Riemannian symmetric spaces of rank one*, Lecture Notes in Pure and Applied Mathematics, Vol. 5. Marcel Dekker, Inc., New York, 1972. vii+81 pp, MR 0383304, Zbl 0239.53032.
- [5] T. Chu, *The Weil-Petersson metric in the moduli space*, Chinese J. Math. **4** (1976), no. 2, 29–51, MR 0590105, Zbl 0344.32006.
- [6] K. Corlette, *Archimedean superrigidity and hyperbolic geometry*, Ann. of Math. (2) **135** (1992), no. 1, 165–182, MR 1147961, Zbl 0768.53025.
- [7] G. Daskalopoulos and R. Wentworth, *Classification of Weil-Petersson isometries*, Amer. J. Math. **125** (2003), no. 4, 941–975, MR 1993745, Zbl 1043.32007.
- [8] G. Daskalopoulos and R. Wentworth, *Harmonic maps and Teichmüller theory*, Handbook of Teichmüller theory. Vol. I, 33–109, IRMA Lect. Math. Theor. Phys., 11, Eur. Math. Soc., Zürich, 2007, MR 2349668, Zbl 1161.30032.
- [9] M. Gromov and R. Schoen, *Harmonic maps into singular spaces and p -adic superrigidity for lattices in group of rank one*, Inst. Hautes Études Sci. Publ. Math. No. 76 (1992), 165–246, MR 1215595, Zbl 0896.58024.
- [10] Z. Huang, *Asymptotic flatness of the Weil-Petersson metric on Teichmüller space*, Geom. Dedicata **110** (2005), 81–102, MR 2136021, Zbl 1069.32004.
- [11] Z. Huang, *The Weil-Petersson geometry on the thick part of the moduli space of Riemann surfaces*, Proc. Amer. Math. Soc. **135** (2007), no. 10, 3309–3316 (electronic), MR 2322763, Zbl 1125.30037.
- [12] J. Jost, *Nonlinear methods in Riemannian and Kählerian geometry*, second edition. DMV Seminar, 10. Birkhäuser Verlag, Basel, 1991, 154 pp, ISBN: 3-7643-2685-9 MR. 1138210, Zbl 0732.53003.
- [13] J. Jost and S. T. Yau, *Harmonic maps and superrigidity*, Tsing Hua lectures on geometry & analysis (Hsinchu, 1990–1991), 213–246. Int. Press, Cambridge, MA, 1997. MR 1482040, Zbl 0888.53036.
- [14] S. Kerckhoff, *The Nielsen realization problem*, Ann. of Math. (2) **117** (1983), no. 2, 235–265, MR 0690845, Zbl 0528.57008.
- [15] K. Liu, X. Sun and S. T. Yau, *Good Geometry on the Curve Moduli*, Publ. Res. Inst. Math. Sci. **44** (2008), no. 2, 699–724, MR 2426362, Zbl 1219.14012.
- [16] K. Liu, X. Sun and S. T. Yau, *Canonical metrics on the moduli space of Riemann surfaces*, I. J. Differential Geom. **68** (2004), no. 3, 571–637, MR 2144543, Zbl 1078.30038.
- [17] K. Liu, X. Sun and S. T. Yau, *Canonical metrics on the moduli space of Riemann surfaces*, II. J. Differential Geom. **69** (2005), no. 1, 163–216, MR 2169586, Zbl 1086.32011.
- [18] C. McMullen, *The moduli space of Riemann surfaces is Kähler hyperbolic*, Ann. of Math. (2) **151** (2000), no. 1, 327–357, MR 1745010, Zbl 0988.32012.
- [19] N. Mok, Y. T. Siu and S. K. Yeung, *Geometric superrigidity*, Invent. Math. **113** (1993), no. 1, 57–83, MR 1223224, Zbl 0808.53043.
- [20] W. Roelcke, *Das Eigenwertproblem der automorphen Formen in der hyperbolischen Ebene*, I, II (German), Math. Ann. **167** (1966), 292–337; *ibid.* **168** (1966) 261–324, MR 0243062, Zbl 0152.07705.
- [21] G. Schumacher, *Harmonic maps of the moduli space of compact Riemann surfaces*, Math. Ann. **275** (1986), no. 3, 455–466, MR 0858289, Zbl 0601.32025.

- [22] Y. T. Siu, *The complex-analyticity of harmonic maps and strong rigidity of compact Kähler manifolds*, Ann. of Math. (2) **112** (1980), no. 1, 73–111, MR 0584075, Zbl 0517.53058.
- [23] L. P. Teo, *The Weil-Petersson geometry of the moduli space of Riemann surfaces*, Proc. Amer. Math. Soc. **137** (2009), no. 2, 541–552, MR 2448574, Zbl 1180.30050.
- [24] A. J. Tromba, *On a natural algebraic affine connection on the space of almost complex structures and the curvature of Teichmüller space with respect to its Weil-Petersson metric*, Manuscripta Math. **56** (1986), no. 4, 475–497, MR 0860734, Zbl 0606.32014.
- [25] M. Wolf, *The Teichmüller theory of harmonic maps*, J. Differential Geom. **29** (1989), no. 2, 449–479, MR 0982185, Zbl 0655.58009.
- [26] S. Wolpert, *Noncompleteness of the Weil-Petersson metric for Teichmüller space*, Pacific J. Math. **61** (1975), no. 2, 573–577, MR 0422692, Zbl 0327.32009.
- [27] S. Wolpert, *Chern forms and the Riemann tensor for the moduli space of curves*, Invent. Math. **85** (1986), no. 1, 119–145, MR 0842050, Zbl 0595.32031.
- [28] S. Wolpert, *Geometry of the Weil-Petersson completion of Teichmüller space*, Surveys in differential geometry, Vol. VIII (Boston, MA, 2002), 357–393. Int. Press, Somerville, MA, 2003. MR 2039996, Zbl 1049.32020.
- [29] S. Wolpert, *Families of Riemann surfaces and Weil-Petersson geometry*, CBMS Regional Conference Series in Mathematics, 113. Published for the Conference Board of the Mathematical Sciences, Washington, DC. American Mathematical Society, Providence, RI, 2010, viii+118 pp, ISBN: 978-0-8218-4986-6. MR 2641916, Zbl 1198.30049.
- [30] S. Wolpert, *Geodesic-length functions and the Weil-Petersson curvature tensor*, J. Differential Geom. **91** (2012), no. 2, 321–359, MR 2971291, Zbl 1252.32022.
- [31] S. Yamada, *On the geometry of Weil-Petersson completion of Teichmüller spaces*, Math. Res. Lett. **11** (2004), no. 2-3, 327–344, MR 2067477, Zbl 1060.32005.
- [32] S. K. Yeung, *Representations of semisimple lattices in mapping class groups*, Int. Math. Res. Not. 2003, no. 31, 1677–1686, MR 1981481, Zbl 1037.57011.

DEPARTMENT OF MATHEMATICS, RICE UNIVERSITY, HOUSTON, TEXAS, 77005-1892,
E-mail address: yw22@rice.edu